



IYCN USAID's Infant  
& Young Child  
Nutrition Project



Photo: Aurelio Ayala III


# National interpretations

Responding to the 2009 WHO guidelines in  
Nigeria and other countries

Wasiu Afolabi  
Infant & Young Child Nutrition (IYCN) Project

This presentation was produced through support provided to the Infant & Young Child Nutrition (IYCN) Project by the U.S. Agency for International Development, under the terms of Cooperative Agreement No. GPO-A-00-06-00008-00. The opinions herein are those of the author(s) and do not necessarily reflect the views of the U.S. Agency for International Development.

# USAID's Infant & Young Child Nutrition Project



# Nutrition and HIV in Nigeria

- HIV prevalence among pregnant women: 4.6%
- Exclusive breastfeeding under 6 months: 13.1%
- No strong recommendation on exclusive breastfeeding.
- Still in the process of implementing the 2006 WHO guidelines.



Photo: Oluseyi Akintola

## Reaching a consensus: May 7, 2010

HIV-infected mothers should exclusively breastfeed their infants for the first 6 months of life, introduce complementary feeds at 6 months and continue breastfeeding for 12 months.

Photo: PATH/Evelyn Hockstein




# PMTCT and nutrition stakeholders

- National HIV/AIDS and Sexually Transmitted Disease Control Programme (NASCP)
- National Nutrition Unit
- National PMTCT Task Team
- UNICEF, World Health Organization, USAID
- Network of People Living with HIV/AIDS in Nigeria
- Pediatric Association of Nigeria
- Infant & Young Child Nutrition Project

## How did stakeholders work together?


- Held series of PMTCT Task Team and Nutrition Division meetings from **January to May 2010**.
- Advocated with nutrition community to participate in national dialogue.
- Clarified recommendations and discussed them within the country context.



# Interpreting recommendation 2


- Confusion around recommendation on breastfeeding to 12 months.
- Final consensus: continued breastfeeding to 12 months.

“Strong recommendation. High quality of evidence for first 6 months; low quality of evidence for recommendation re. 12 months.”



# Interpreting recommendation 5

- Confusion around AFASS (acceptable, feasible, affordable, sustainable, and safe) versus one country feeding option.
- Final consensus: Nigeria should adopt one message supporting breastfeeding.





# Zambia response

- National process led by PMTCT.
- At first, recommended no more than 12 months breastfeeding.
- Advocacy from nutrition community led to national 24 month recommendation.
- Concern remains about criteria for stopping breastfeeding at 12 months.



Photo: Kali Erickson

# Lessons learned

- Don't rush the process.
- Encourage nutrition colleagues to engage in national dialogue.
- Bring nutrition and PMTCT stakeholders together early on.
- Build a broad understanding of the recommendations among stakeholders.
- Consider the country context.



Photo: Oluseyi Akintola

## What's next?

- Finalize the national consensus statement.
- Develop national communication strategy.
- Incorporate recommendations into national guidelines and training curricula.
- Implement in facilities and communities.



Photo: Oluseyi Akintola

Thank you



Photo: David and Lucile Packard Foundation